

blank page

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Apparent 

Phosphite 600

SYSTEMIC FUNGICIDE

ACTIVE CONSTITUENT: 600 g/L PHOSPHOROUS (PHOSPHONIC) ACID
PRESENT AS MONO (AND) DI POTASSIUM PHOSPHITE

GROUP 33 FUNGICIDE

Effective against Phytophthora diseases and Downy Mildew in various situations as per Directions for Use Table.

IMPORTANT: Read this booklet before use.

APVMA Approval No: 68156/57644

AIRR APPARENT PTY LTD
15/16 Princes Street, Newport NSW 2106
P 03 5820 8400 | E enquiries@apparentag.com.au | www.apparentag.com.au

DIRECTIONS FOR USE

Restraint: DO NOT apply this product at volumes which cause excessive run off.

TREE AND VINE CROPS

CROP	DISEASE	STATE	TREATMENT METHOD	RATE
AVOCADO	Phytophthora Root Rot (<i>Phytophthora cinnamomi</i>)	QLD, NSW, SA, VIC, WA, ACT	Injection	Trunk Injection, Skeletal trees: 1st year 5 mL undiluted product per metre of canopy diameter. Other situations: 2.5 mL product diluted with 7.5 mL water per metre of canopy diameter.
			Foliar Spray	3.3 – 4 L/ha
YOUNG OR SMALL CITRUS	Phytophthora Root Rot (<i>P. nicotianae</i>), and Collar Rot (<i>P. citrophthora</i>)	All states	Foliar spray	170 - 330 mL/ 100 L to leaf wetness (by boom or similar high volume sprayer)
MATURE CITRUS		All states	Foliar spray	13.3 L/ha in 3000 - 4000 L of water. 53 mL/12 L per tree. 8.3 L/ha in 2000 - 5000 L of water. 33 mL/12 L per tree
GRAPES	Downy Mildew (<i>Plasmopara viticola</i>)	All States	Foliar spray	2 L/ha early season, small canopies 2.7 - 4 L/ha Mid-late season, large canopies

CRITICAL COMMENTS
<p>Inject trees at spring flush maturity - Repeat treatment in February or March. Drill holes 5 mm in diameter and 25 – 50 mm deep with slight downward angle in trunk. Syringes should be placed in the main trunk of the tree and spaced evenly around the circumference of the trunk. Suitable for use with Chemjet tree injectors, Ag-murf gun or hydraulic tree injection. Do not prune trees back before injection process as burning of new growth may occur. Do not inject trees in winter months. Do not cut back the canopy of injected trees. Do not add anything other than water to Apparent Phosphite 600 for trunk injection. DO NOT inject more liquid in a lesser number of syringes than directed.</p> <p>Solution concentrate: 165 - 200 mL/100 L. Amount per unit: Spray to run off – 7.5 to 10 L per adult tree.</p> <p>1. Curative: Apply every 3 weeks until disease is under control. 2. Preventative: Apply every 5 - 6 weeks.</p> <p>Two applications <i>1st Application:</i> Late winter prior to flowering. <i>2nd Application:</i> Autumn applied to mature fruit. Repeat applications annually to maintain protection for the plant. Depending on condition at time of application use a non-ionic wetting agent in conjunction with Apparent Phosphite 600. WHERE DISEASE INCIDENCE IS HIGH OR WELL ESTABLISHED. For effective control, apply as a protectant before above ground symptoms of decline and collar rot become evident, spray trees for even coverage. Do not apply under high temperatures (above 35°C) particularly if humidity is low, or to moisture stressed trees. LOW PHYTOPHTHORA PRESSURE, WELL DRAINED SOIL Removal of fruit from affected plants will enhance recovery. WARNING. Soft skin citrus trees may develop leaf burn and growth retardation following foliar application or soil drench of Potassium Phosphonate at rates recommended for established trees. WARNING: 1. Foliar application to fruit may cause damage to fruit. 2. Soft skinned citrus trees may develop leaf burn and growth retardation following foliar application or soil drench.</p> <p>Before applying this product for the first time, a small test area should be sprayed and observed for at least 1 week.</p> <p>It is essential that the rate of Apparent Phosphite 600 is adjusted to the vine-row volume (i.e. the volume of vine foliage per hectare). An application volume of 500 L/ha is suggested at the start of the season, increasing to 2000 L/ha in a vigorous crop at full canopy. Spray timing is critical. For best results apply Apparent Phosphite 600 as a tank mix with protectant fungicides such as Mancozeb/Dithane, Copper Oxchloride etc, to ensure both pre- and post-infection activity. Apparent Phosphite 600 should be applied at times of high disease risk, especially between the time that conditions are conducive to Downy Mildew infection and the appearance of oil spots. Ensure spray coverage is adequate and that the appropriate rate of Apparent Phosphite 600 is applied to match vine growth, particularly from mid-season onwards and especially where grapes are grown on root stock.</p>

NON TREE AND VINE CROPS

CROP	DISEASE	STATE	TREATMENT METHOD
CUCURBITS	Downy Mildew (<i>Pseudoperonospora cubensis</i>)	All States	Foliar spray
ORNAMENTALS	Phytophthora Root and Crown rot (<i>Phytophthora</i> spp)	Qld, NSW, Vic, WA, TAS, NT, ACT	Foliar spray
PINEAPPLES	Phytophthora Root (<i>Phytophthora cinnamomi</i>) and Heart Rot (<i>Phytophthora parasitica</i>)	Qld, WA	Foliar spray
SUBTERRANEAN CLOVER	Phytophthora Root Rot (<i>Phytophthora clandestina</i>)	NSW, SA VIC, WA, TAS, ACT	Foliar spray

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

Subterranean Clover: **DO NOT GRAZE OR CUT FOR STOCKFEED FOR 14 DAYS AFTER APPLICATION.**

Citrus, Cucurbits, Avocado, Grapes, Pineapples: **NOT REQUIRED WHEN USED AS DIRECTED.**

GENERAL INSTRUCTIONS

For effective disease control a good leaf cover must be achieved before a widespread disease outbreak occurs.

CONDITIONS CONDUCIVE TO DOWNY MILDEW INFECTION

(1) PRIMARY INFECTION - overnight conditions of:

- Temperature 10°C
- Rainfall 10 mm
- Soil Wetness 24 hrs
- Leaf Wetness 3 - 4 hrs at end of a 24 hr period.

(2) SECONDARY INFECTION - Overnight conditions of:

- Temperature 11°C
- Humidity 98% for at least 4 hours from midnight to dawn
- Leaf Wetness 24 hrs

TO AVOID RESISTANT STRAINS OF DOWNY MILDEW DEVELOPING, APPARENT PHOSPHITE 600 SHOULD BE APPLIED AS CLOSE AS POSSIBLE TO THE DAY OF INFECTION AND ALTERNATED WITH SYSTEMIC FUNGICIDES.

RATE	CRITICAL COMMENTS
3 L /ha in min 800 - 1000 L of water	Use weekly spray intervals when conditions favour disease development. To avoid phytotoxicity with some plant species, it is recommended that the product be tested on a few plants of each species prior to the main application.
Knapsack/Boom 170 mL/ 100L Air Blast 330 mL/100 L	Apply at 4 - 6 weekly intervals when conditions favour disease development. To avoid phytotoxicity with some plant species it is recommended that the product be tested on a few plants of each species prior to the main application. DO NOT apply when ornamental plants are dormant or stressed. DO NOT apply to ornamental plants under extremes of temperature.
4 L/ha 1:200 - 1:500 water	Apply to tops, two (2) weeks prior to harvest of planting material.
500 mL/ha approx. 200 L water/ha	Apply 8 to 9 days after first irrigation but before second irrigation. Apply in Autumn when Subterranean Clover is at the cotyledon to unifoliate leaf growth stage.

FUNGICIDE RESISTANCE WARNING

GROUP 33 HERBICIDE

For fungicide resistance management, Apparent Phosphite 600 is a Group 33 fungicide.

Some naturally occurring individual fungi resistant to Apparent Phosphite 600 Systemic Fungicide and other Group 33 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Apparent Phosphite 600 Systemic Fungicide, and other Group 33 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, AIRR Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Phosphite 600 Systemic Fungicide to control resistant fungi.

MIXING

Half fill spray vat and with agitation system running, add the required amount of product.

COMPATIBILITY

Compatible with Mancozeb, Copper Oxchloride, Bayleton, Tilt, and most common powdery mildew fungicides. Most foliage nutrients are also compatible. When using in conjunction with 'EC's' always form emulsion with water prior to adding Apparent Phosphite 600 Systemic Fungicide.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

This product is harmful to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Spillage should not be directed to drains but absorbed in sawdust and the absorbent material disposed of in a sealed container at an approved disposal site. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If not recycling, replace cap and return clean containers to recycler or designated collection point.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to collection point for refill or storage.

If not recycling break, crush or puncture containers and bury empty containers in a local authority landfill. If no landfill is available, bury containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

May irritate eyes and skin. Avoid contact with eyes and skin. Do not inhale spray mist. When preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves, goggles and disposable face mask. If clothing becomes contaminated with product or wet with spray, remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which is available from the supplier.

CONDITIONS OF SALE

The use of Apparent Phosphite 600 Systemic Fungicide being beyond the control of the manufacturer no warranty expressed or implied is given by AIRR Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AIRR Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.