

blank page

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



MCPA 750

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 750 g/L MCPA
present as the DIMETHYLAMINE SALT

GROUP | **HERBICIDE**

For the control of broadleaf weeds in cereals, linseed, pastures, rice, sugar cane and turf as specified in the Directions for Use table.

IMPORTANT: Read the attached booklet before use.

APVMA Approval No. 68355/58213

AIRR APPARENT PTY LTD
15/16 Princes Street, Newport NSW 2106
P 03 5820 8400 | E enquiries@apparentag.com.au | www.apparentag.com.au

DIRECTIONS FOR USE

Restrictions: DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply if rain is likely within 6 hours.

1. FIELD CROPS

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha
Wheat, Barley, Oats, Triticale	Refer to Weed Table	TAS, SA only	460 mL - 1.45 L
		QLD only	
		NSW, ACT only	460 – 960 mL
		VIC only	
Wheat, Barley, Triticale, Cereal Rye	Capeweed, Double Gee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish	WA only	600 mL - 1.35 L
		WA only	270 mL plus 350 mL diuron (500 g/L flowable)
	Amsinckia (Yellow Burr), Capeweed, Charlock Deadnettle, Double Gee (Spiny Emex, Three Corner Jack), <i>Erodium</i> sp., Mustard, Turnip Weed, White Ironweed (Corn Gromwell, Sheep Weed), Wild Radish, Wild Turnip	NSW, ACT, VIC only	330 mL plus 500 mL diuron (500 g/L flowable)
		SA only	230 mL plus 500 mL diuron (500 g/L flowable)
Wheat, Barley, Oats, Triticale	Soldier Thistle	SA only	460 mL plus 200 mL Apparent Clopyralid 300
	Skeleton Weed		660 mL plus 500 mL Apparent Clopyralid 300
Field Peas	Prickly Lettuce (<i>Lactuca serriola</i>)	NSW, VIC, TAS, SA only	Weed Stage: 4 leaf, 80 mL plus 125 mL diflufenican (500 g/L)
			Weed Stage: 6 leaf, 100 mL plus 150 mL diflufenican (500 g/L)
	Charlock (Wild Mustard) (<i>Sinapis arvensis</i>)	NSW, VIC, TAS, SA only	Weed Stage: 4 leaf, 80 mL plus 125 mL diflufenican (500 g/L)
			Weed Stage: 6 leaf, 100 mL plus 150 mL diflufenican (500 g/L)

CRITICAL COMMENTS

Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing.

Apply from early tillering (4 to 5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem).

Apply 460 mL rate after crop has reached 5 leaf stage and weeds marked in weed table have 3 - 6 leaves. Higher rates after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.

Apply 460 mL rate after crop has reached 2-3 leaf stage and weeds marked # in weed table have 2 - 4 leaves. Higher rate after the crop plants have 5 leaves to fully tillered.

Apply to crop from 5 leaf to Flag leaf emergence (Zadoks 15 - 37).

CROP: 3 - 4 leaf stage (Zadoks 13 - 14).

CROP: 4 - 5 leaf stage (Zadoks 14 - 15).

Apply by aircraft or boom with 20 – 100 L/ha water, when most weeds have germinated and are in 2 - 5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application.

Undersown sub-clovers may be slightly retarded.

DO NOT apply to undersown medics.

Apply by boom spray or aircraft when the crop is at the 3 - 5 leaf stage (NSW, ACT and SA only) and 5 leaf stage (VIC only).

Weeds must be in the 2 - 5 leaf stage.

DO NOT apply to undersown medic or lucerne.

DO NOT use on Kamilaroi, Kite, Olympic or Shortim Wheat.

DO NOT use on very light sandy soils.

Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing.

Apply to rosettes from 5 - 15 cm diameter when the crop is between the leaf and late tillering growth stages. It is essential that all the Skeleton weed is emerged and minimum of 5 cm diameter which is usually not before August. This rate will provide control of Skeleton Weed until harvest and will substantially reduce seed numbers the following season.

FIELD PEAS: Apply early post-emergence after the third node stage and before the start of flowering.

Warning: Field peas grown on high pH soils in the presence of free lime may be less tolerant to diflufenican (500 g/L).

APPLICATION AND WEED CONTROL: Apply when weeds are actively growing. For optimum results apply 4 to 6 weeks post-sowing.

Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as:

- high crop and weed density,
- late season germinations,
- abnormal weed growth (including early flowering); higher rates of product (up to the maximum rate of application specified for that weed) may be required.

Diflufenican (500 g/L) will not effectively control:

- regrowth of suppressed weeds,
- transplanted weeds,
- regrowth from rhizomes or roots,
- weeds growing under stress from previous herbicide applications.

1. FIELD CROPS continued

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha
Linseed	Refer to Weed Table	QLD only	460 mL - 1.45 L
		NSW, ACT, VIC, SA, TAS only	460 – 730 mL
Sugar Cane	Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory As above plus Fat Hen, Noogoora Burr	QLD only	930 mL
			1.45 L
Rice	Dirty Dora, Starfruit, Sedges, Buddah (Butter) Pea	NSW only	485 mL to 970 mL
Flax	Refer to Weed Table	VIC, TAS only	325 mL

2. PASTURES, SEED CROPS & TURF

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha	
Grass Pastures	Refer to Weed Table	All States	460 mL - 2.7 L	
	Salvation Jane	SA only	660 mL plus 400 mL diuron (500 g/L flowable)	
	Soldier Thistle		460 mL - 1.45 L	
	Seedling and established Horehound	NSW, ACT only	2 L	
Grass Seed Crops	Refer to Weed Table	QLD, VIC, TAS, SA, WA only	460 mL-2.7 L	
Sub-Clover Seed Crops	Refer to Weed Table	VIC, SA only	460 – 960 mL	
	Pastures – Spray/Graze Technique	Calltrop, Capeweed, Charlock, Mustards, Paterson's Curse, Shepherd's Purse, Saffron, Slender Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	NSW, ACT, VIC only	230 – 930 mL
		Refer to Weed Table	TAS only	460 mL
	Capeweed, Dandelion, Seedling Dock, Three Corner Jack, Geranium, Erodium, Mustard, Annual Thistles, Turnip Weed, Wild Radish, Wild Turnip Soldier Thistle	WA only SA only	610 mL - 1.25 L 330 mL	
Turf	Refer to Weed Table	All States	330 mL or 2.8 mL per 100 sq. metres	
			930 mL - 1.8 L or 10 – 19 mL per 100 sq. metres	

3. SPOT SPRAYING

SITUATION & CROP	WEEDS CONTROLLED	STATE
High Volume Spraying	Refer to Weed Table	All States
Knapsack Application	Refer to Weed Table	All States

CRITICAL COMMENTS
Apply when crop is 10 – 15 cm tall with at least 170 L/ha of water. DO NOT spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. DO NOT apply by aircraft.
Post-emergent: Apply as a directed inter-row spray. DO NOT exceed rates with phenoxy sensitive varieties.
Apply by aircraft at early/mid tillering stage of rice but not later than panicle initiation. Use 485 mL/ha at 35 days after aerial sowing (approx. 2 litres per rice plant). Use 970 L/ha at 45 days or more after cereal sowing (3 tillers or more per rice plant). The crop should be sprayed before flowering of the weeds. Water Management: Before herbicide application, lower water level in the bays to at least two thirds of the weed growth for direct contact with MCPA 750. Weeds covered by water will not be controlled. Raise water depth to normal levels 24 hours following spraying. Apply to crop when 8 – 20 cm high.

CRITICAL COMMENTS
Apply to established grass based pastures. Damage may occur to legumes if present. See "Legume Tolerance" below. NOTE: Capeweed, Paterson's Curse and Variegated Thistle may be poisonous to stock after spraying. Use this mixture only if spraying is delayed until early winter. Rosettes should not exceed 20 cm diameter.
Use low rate on seedlings with no more than 2 true leaves. Use 100 litres of water for best results. If clover present do not exceed 575 mL. Apply in Autumn when horehound plants are at the seedling stage. If sub-clover or medic plants are present they will be damaged by this application. Apply to established grass seed crops.
DO NOT exceed rate specified. Spray in Autumn and early Winter. Spray actively growing 6 - 8 weeks old weeds. Introduce stock 7 - 10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and medic may be damaged and should be grazed short before spraying. See "Legume Tolerance" under General Instructions. Precaution: An increased quantity of poisonous plants may be eaten by stock using spray/graze e.g. Calltrop, Capeweed, Paterson's Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson's Curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. DO NOT graze horses or pigs on Paterson's Curse. Observe withholding period of at least 7 days after spraying. Newly laid: Delay until well established or at least two months old. Apply in high volume to actively growing weeds. See "Legume Tolerance". Buffalo Grass is more easily damaged than other grasses. Established: Apply in high volume to actively growing weeds. DO NOT mow for 2 days before application. Some transitory damage may occur to fine turf grasses and strawberry clover. See "Legume Tolerance". Buffalo Grass is more easily damaged than other grasses.

MIXING RATE/COMMENTS
Use 130 mL/100 L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. Use 270 mL/100 L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed. Use 13 mL/10 L of water for easy to kill weeds such as Wild Turnip, Mustards and Wild Radish. Use 27 mL/10 L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed.

4. NON-CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY

SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha
Non-Crop areas, Fallow Land, Industrial and Commercial Areas, Rights-of-way	Refer to Weed Table	All States	Refer to Weed Table

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

CEREALS, PASTURE: DO NOT GRAZE OR CUT STOCK FOOD FOR 7 DAYS AFTER APPLICATION. HARVEST WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

WEED TABLE:

Where weeds are to be sprayed in a crop or pasture situation, use only the rate given for the particular crop or situation indicated under the Directions for Use table. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those States where a rate or range of rates is indicated for the particular weeds listed.

APPLICATION RATE PER HECTARE

* Refer to Critical Comments for Wheat, Barley, Oats, Triticale

WEEDS	APPLICATION RATE (per hectare)					
	VIC	NSW, ACT	SA	TAS	QLD	WA
Ball Mustard	-	-	1.15 L	-	-	-
Bathurst Burr	1.35 L	660 mL - 1.35 L	1.35 L	1.35 L	1.35 L	1.35 L
Black Bindweed	-	970 mL - 1.35 L	-	-	930 mL - 1.35 L	-
Californian Burr	-	660 mL - 1.35 L	-	1.45 L	-	-
Cape Tulip	-	-	-	-	-	1.35 L
Capeweed	-	1.45 L	1.45 L	1.45 L	1.45 L	-
Carrot Weed	1.35 L	-	-	-	1.35 L	-
Charlock	460 mL	660 mL	460 mL	660mL - 1.45L	460 mL	-
Cutleaf Mignonette	-	-	1.45 L	-	-	-
Dandelion	1.35 L	1.35 L	1.35 L	1.35 L	1.35 L	1.35 L
Deadnettle	-	1.45 L	-	-	1.45 L	-
Docks	960 mL	-	1.45 L	1.45 L	-	-
Fat Hen	1.35 L	660 mL - 1.35 L	-	1.35 L	1.35 L	-
Fennel	1.8 L	1.8 L	1.8 L	1.8 L	1.8 L	-
Field Bindweed	1.35 L	-	1.35 L	1.35 L	1.35 L	1.35 L
Field Cress	-	-	-	1.45L	-	-
Flatweed or Catsear	-	-	-	1.45 L	-	930 mL
Fumitory - red	930 mL	930 mL	930 mL	-	930 mL	-
Hedge Mustard	730 mL*	-	460 mL	460 mL	460 mL	825 mL
Hexham Scent or Melilotus	960 mL	800 mL	930 mL	-	930 mL	-
Hoary Cress	3 L	-	1.45 L	-	3 L	-
Horehound	-	2 L	1.45 L	-	-	-
Lesser Swinecress	-	-	-	1.45 L	-	-
Lincoln Weed	-	-	460 mL	-	-	-
London Rocket	-	-	-	-	-	825 mL

CRITICAL COMMENTS
Apply at rates shown in Weed Table when weeds only are present and non crop or pasture is involved.

CRITICAL COMMENTS
Apply at rosette stage.
Spray young seedlings only.
Apply at young stage prior to flowering.
Spray young seedlings only.
Apply just prior to flowering. Gives mild suppression only.
Apply at seedling stage.
Apply at seedling stage.
Spray up to rosette stage.
Apply at young rosette stage.
Apply at young rosette stage.
Moderately susceptible. Apply at seedling stage.
Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected.
Spray up to rosette stage.
Apply at young stage.
Apply at bud stage, repeating over several seasons.
Apply at young stage.
Apply when weeds are 10 - 15 cm high.
Apply at young stage prior to flowering.
Apply at rosette stage.
Apply at young seedling stage.
Apply before flowering.
Apply at rosette stage.
Apply at young stage.
Apply at rosette stage.

APPLICATION RATE PER HECTARE continued

WEEDS	APPLICATION RATE (per hectare)					
	VIC	NSW, ACT	SA	TAS	QLD	WA
Lupins	-	460 mL - 960 mL	-	-	-	825 mL
Mintweed	1.35 L	1.35 L	-	-	1.35 L	-
Mustards	730 mL*	660 mL	730 mL	730 mL	-	825 mL
Neogoorra Burr	1.35 L	660 mL - 1.35 L	1.35 L	-	1.35 L	1.35 L
Opium Poppy	-	-	-	1.45L	-	-
Paterson's Curse	-	660 mL - 960 mL	-	-	2.7 L	-
Pimpernel	-	-	460 mL	-	-	-
Plantains	-	1.35 L	-	1.35 L	1.35 L	-
Rapeseed	-	660 mL - 960 mL	-	-	-	-
Rough Poppy	-	460 mL - 960 mL	-	-	-	-
Safflower	-	460 mL - 660 mL	-	-	-	-
Shepherds Purse	-	-	-	1.45 L	-	-
Skeleton Weed	-	960 mL - 1.35 L	-	-	-	-
Sorrel	-	-	1.45 L	-	-	-
Stinkwort	1.35 L	660 mL - 960 mL	1.35 L	-	-	1.35 L
Sunflower (seedlings)	-	690 mL - 1 L	-	-	-	-
Thistles - Artichoke	1.05 L	-	1.45 L	-	1.35 L	460 mL
- Nodding	-	1.65 L	-	-	-	-
- Saffron	960 mL - 1.65 L	660 mL - 1.35 L	930 mL - 1.65 L	930 mL - 1.65 L	930 mL - 1.65 L	960 mL - 1.65 L
- Scotch or Cotton	1.45 L	-	460 mL - 930 mL	1.45 L	-	-
- Slender	2 L	660 mL - 1.35 L	660 mL	2 L	-	2 L
- Soldier	730 mL	-	730 mL	-	-	-
Thistles - Spear	-	960 mL - 1.35 L	-	1.45 L	-	-
- Star	-	-	730 - 930 mL	-	-	-
- Variegated	960 mL	660 mL - 1.35 L	460 mL - 930 mL	960 mL	960 mL	960 mL
Turnip Weed	730 mL*	660 mL*	660 mL	-	-	-
Wild Artichoke	-	-	1.45 L	-	-	-
Wild Radish	930 mL	660 mL*	930 mL	1.35 L	460 mL - 730 mL	800 mL
Wild Turnip	730 mL	660 mL*	460 mL	1.35 L	460 mL - 730 mL	600 mL

GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours following application.

DO NOT spray crops or weeds outside the stages indicated in 'Critical Comments' as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

APPLICATION/MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

BOOM SPRAYING

- Use 30 - 120 litres of water per hectare.
- Use minimum of 170 litres water per hectare for LINSEED.

AERIAL SPRAYING

- Use 10 - 90 litres water per hectare.

EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with Spraymate Tank and Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

CRITICAL COMMENTS

Spray up to 10 cm high.
 Spray young seedlings only.
 Spray up to rosette stage.
 Spray young seedlings only.
 Apply at young stage.
 Apply at early rosette stage.
 Apply at rosette stage.
 Apply at young stage.
 Spray up to rosette stage.
 Spray up to rosette stage.
 Spray up to rosette stage.
 Apply at rosette stage.
 Spray rosettes before aerial growth commences.
 Apply at rosette stage.
 Apply when weeds are 10 - 15 cm high.
 Spray up to rosette stage.
 Apply at rosette stage.
 Spray early rosettes. Repeat spray required in following Spring or Autumn.
 Spray up to rosette stage (September WA).
 Apply at seedling stage.
 Spray young seedlings only.
 Spray young seedlings. Use higher rate for larger weeds.
 Apply at rosette stage.
 Spray young seedlings. Use higher rate for larger weeds.
 Spray young seedlings only. Use 1.35 L/ha (SA) for well developed weeds.
 Spray up to rosette stage.
 Apply at rosette stage.
 Spray up to rosette stage.
 Spray up to rosette stage.

RESISTANT WEEDS WARNING

GROUP I	HERBICIDE
----------------	------------------

Apparent MCPA 750 Selective Herbicide is a member of the Phenoxy group of herbicides. This product has the disruptors of plant cell growth mode of action. For weed resistance management this product is a Group I Herbicide. Some naturally-occurring weed biotypes resistant to this product and other Group I herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group I herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, AIRR Apparent Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

COMPATIBILITY/TANK MIXES

This product can be tank mixed with Apparent Metsulfuron 600, Banvel[®], Apparent Bromoxynil 200, Flowable Diuron, Apparent Clopyralid 300, Sertin[®], Apparent Paraquat 250, Flowable Terbutryn, Tordon[®] and Tribunil[®]. The effectiveness of Imidan[®], Le Mat[®], Metasystox[®] and dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of Spraybuff[®] or Spraymate[®] L1-700 is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly, before spraying.

PROTECTION OF CROPS, NATIVE AND OTHER NON TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto healthy susceptible plants/crops, cropping lands or pastures.

Drift Warning: DO NOT use on or in situations where damage to susceptible crops – such as cotton, vines, tomatoes, vegetables, fruit trees, lucerne, legume crops and pastures, oilseed crops or other susceptible crops and trees (eg. Kurrajongs, Belahs, Eucalypts) – may occur from direct application, spray drift or vapours.

LEGUME TOLERANCE

The use of MCPA is preferred to 2,4-D Amine (Amicide) or Ester (Estercide) in situations where legumes are undersown in the crop.

Note: The tolerance of pasture legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with lucerne as damage may result.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers or streams with the product or used containers.

STORAGE AND DISPOSAL

20 L, 200 L: Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

110 L: Store in the original sealed Envirodrum in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the drum with water or any other foreign matter. After each use of the product, please ensure the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drums remain the property of AIRR Apparent Pty Ltd.

1000 L: Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs AIRR Apparent Pty Ltd should be notified immediately. This minibulk/bulk container is reusable and remains the property of AIRR Apparent Pty Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to AIRR Apparent Pty Ltd for cleaning, relabelling and refilling.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes. Will irritate the skin. Avoid contact with the eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product on skin, immediately wash area with soap and water. If product or spray in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia 131126.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A material safety data sheet for Apparent MCPA 750 Selective Herbicide is available from AIRR Apparent Pty Ltd upon request.

CONDITIONS OF SALE

The use of Apparent MCPA 750 Selective Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by AIRR Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AIRR Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

[^]Other trademarks