

blank page

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Apparent 

Poacher 750

HERBICIDE

ACTIVE CONSTITUENT: 750 g/kg IMAZAPYR

GROUP B HERBICIDE

For the control of various annual and perennial weeds in non-crop situations as indicated in the Directions for Use table.

IMPORTANT: Read this booklet before use.

APVMA Approval No: 69538/102235

AIRR APPARENT PTY LTD
15/16 Princes Street, Newport NSW 2106
P 03 5820 8400 | E enquiries@apparentag.com.au | www.apparentag.com.au

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply by AIRCRAFT as spray drift may cause damage to desirable surrounding species.

Situation	Weed
<p>Around agricultural buildings and other farm non-crop situations, commercial, industrial, and public service areas, rights of way and waste land, away from desirable vegetation.</p>	<p>Annual ryegrass (<i>Lolium rigidum</i>), Barnyard grass (<i>Echinochloa crus-galli</i>), Blady grass (<i>Imperata cylindrica</i>), Buckbush (<i>Salsola kali</i>), Deadnettle (<i>Lamium amplexicaule</i>), Feathertop Rhodes grass (<i>Chloris virgata</i>), Fleabane (<i>Coryza canadensis</i>), Hexham Scent (<i>Melilotus indicies</i>), Liverseed grass (<i>Urochloapanicoides</i>), London rocket (<i>Sisymbrium irio</i>), Mayne's pest (<i>Verbena tenuiseecta</i>), Medics (<i>Medicago</i> spp.), Mintweed (<i>Salvia reflexa</i>), Blackberry Nightshade (<i>Solanum nigrum</i>), Paterson's curse (<i>Echium plantagineum</i>), Plains grass (<i>Stipa aristiglumis</i>), Prairie grass (<i>Bromus unioloides</i>), Prickly Lettuce (<i>Lactuca serriola</i>), Purple Top (<i>Verbena bonariensis</i>), Scented Top (<i>Capillipedium</i> spp.), Sedges (<i>Cyperus</i> spp.), Slender Celery (<i>Apium leptophyllum</i>), Sowthistle (<i>Sonchus oleraceus</i>), St Barnaby's thistle (<i>Centaurea solstitialis</i>), Summer grasses (<i>Digitaria</i> spp.), Thornapples (<i>Datura</i> spp.), Turnip weed (<i>Rapistrum rugosum</i>), Variegated Thistle (<i>Silybum marianum</i>), Vasey grass (<i>Paspalum urvillei</i>), Warrego grass (<i>Paspalidium juborum</i>), Wild Oats (<i>Avena</i> spp.), Wild Turnip (<i>Brassica tournefortii</i>), Wireweed (<i>Polygonum aviculare</i>)</p>
<p>Around agricultural buildings and other farm non-crop situations, commercial, industrial, and public service areas, rights of way and waste land, away from desirable vegetation.</p>	<p>Bokhara clover (<i>Melilotus albus</i>), Bracken (suppression only) (<i>Pteridium esculentum</i>), Capeweed (<i>Arctotheca calendula</i>), Couch (<i>Cynodon dactylon</i>), Docks (<i>Rumex</i> spp.), Johnson grass (<i>Sorghum halepense</i>), Panics (suppression only) (<i>Panicum</i> spp.), Paspalum (<i>Paspalum dilatatum</i>), Queensland bluegrass (<i>Dichanthium sericeum</i>), Stinging Nettle (<i>Urtica incisa</i>)</p>

State	Rate	Critical Comments
All States	1.3 kg/ha	<p>Applications can be made pre- or post-emergence. Post-emergence applications also provide pre-emergence control.</p> <p>Medium Volume: 1.3 kg Apparent Poacher 750 Herbicide should be applied using spray volumes ranging between 50 and 200 L/ha.</p> <p>High Volume: Apply sufficient spray to wet the surfaces visibly without producing runoff. Measure how much spray is required to cover 10 m x 10 m. Add Apparent Poacher 750 Herbicide at the rate of 13 g in this measured volume.</p>
All States	2 kg/ha	<p>Applications can be made pre- or post-emergence. Post-emergence applications also provide pre-emergence control.</p> <p>Medium Volume: 2 kg Apparent 750 SG should be applied using spray volumes ranging between 50 and 200 L/ha.</p> <p>High Volume: Apply sufficient spray to wet the surfaces visibly without producing runoff. Measure how much spray is required to cover 10 m x 10 m. Add Apparent 750 SG at the rate of 20 g in this measured volume.</p>

Situation	Weed
Around agricultural buildings and other farm non-crop situations, commercial, industrial, and public service areas, rights of way and waste land, away from desirable vegetation.	Rubber vine (<i>Cryptostegia grandiflora</i>)
	Thunbergia (<i>Thunbergia</i> spp.)

State	Rate	Critical Comments
Qld only	1.3 g/L water	Post emergence application. Apply sufficient spray to wet the surface visibly to the point of run-off. For effective control, apply when the plant is actively growing and not stressed, yellowing or bearing pods. Flowering is a good indication of the best time for application. Use high volume application as detailed in the General instructions section.
	2.5 g/L water	Post emergence application. Apply sufficient spray to wet the surface visibly to the point of run-off. For effective control, apply when the plant is actively growing. Use high volume application as detailed in the General instructions section.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Apparent Poacher 750 Herbicide is a soluble granule herbicide which will kill a range of standing weeds and emerging seedlings. It stops growth rapidly, although symptoms may take over 30 days to develop. Affected plants are discoloured and often growth is distorted.

The active material in Apparent Poacher 750 Herbicide is rapidly absorbed through foliage and roots, and translocated throughout plants. Foliage treatments will provide pre-emergence control as the active material is released to the soil following plant decay.

Residual control of weeds, for which the lower rate of Apparent Poacher 750 Herbicide is recommended, will increase when the higher rate is used.

RESISTANT WEEDS WARNING

GROUP B HERBICIDE

Apparent Poacher 750 Herbicide is a member of the imidazolinone group of herbicides. Apparent Poacher 750 Herbicide has the inhibition of acetolactate synthase (ALS) mode of action. For weed resistance management, Apparent Poacher 750 Herbicide is a Group B herbicide. Some naturally-occurring weed biotypes resistant to Apparent Poacher 750 Herbicide and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Apparent Poacher 750 Herbicide or the Group B herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, AIRR Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Poacher 750 Herbicide to control resistant weeds.

APPLICATION

Faster control of standing weeds will result if growing conditions are favourable at and after application. Rainfall within two hours of the spray mixture drying on leaves may result in reduced weed control.

SPRAYING INSTRUCTIONS

For medium volume spraying, apply through a spray boom calibrated to ensure even coverage. Use spray nozzles suitable to prevent spray drift onto surrounding desirable vegetation.

For high volume application, use a hand gun or knapsack to apply sufficient spray to wet the surface visibly without producing run-off.

MIXING

Pour the required amount of Apparent Poacher 750 Herbicide into a spray container which already holds the total amount of water required. Mix thoroughly. If Apparent Poacher 750 Herbicide is added during filling, foaming may occur. If excessive foaming is a problem, add a suitable antifoaming agent. Consult your distributor for specific information.

Apparent Poacher 750 Herbicide may be applied in hard or soft water. The product is corrosive to mild steel. Use ONLY stainless steel, fibreglass, plastic or plastic-lined containers for mixing, storage and application.

COMPATIBILITY

For faster knockdown of weeds, certain other herbicides may be mixed with Apparent Poacher 750 Herbicide.

Bipyridyl herbicides (eg. paraquat) are NOT compatible with Apparent Poacher 750 Herbicide. Mixtures of this type will result in poor weed control.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Do NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

Do NOT apply on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots.

Do NOT apply where run-off water may flow onto agricultural land or other desirable vegetated areas. Dispose of washings away from desirable plants and their roots.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves, goggles and disposable dust mask. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a Doctor or Poisons Information Centre. Phone Australia: 13 11 26.

Safety Data Sheet

Additional information is listed in the Safety Data Sheet which is available from the supplier.

CONDITION OF SALE

The use of Apparent Poacher 750 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by AIRR Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AIRR Apparent Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.