SITUATIONS	PEST	RATE (kg/ha)	RATE kg/100 m <sup>2</sup>	CRITICAL COMMENTS
External surrounds of building and structures: Gardens, Lawns, Ant nests & trails (cont)	Fleas (Adults & Larvae) (Ctenocephalides spp.)	110-220	1.1-2.2	Flea larvae develop in the soil of shade areas that are accessible to pets or of their animals sults irrigate the treated areas with up to 10 mm of water immediately after application. Use higher application rates when treating areas with dense ground cover heavy leaf little.
	Ticks (Ixodes spp.)	110-220	1.1-2.2	Applications should be made in late summer/autumn to control language and inymphs that reside in the soil and leaf little DO NOT make soot applications. Ireal the entire area where exposure to ticks may occasing areas with dense ground cover heavy leaf little. Ticks may be entroduced from surrounding areas on host animats.

### NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED.

### GENERAL INSTRUCTIONS

For optimum results use a properly calibrated INSECTICIDE RESISTANCE WARNING

## GROUP 3/A INSECTICIDE

For insecticide resistance management Apparent Antassassin Insecticide is a Group 3A insecticide. Some naturally occurring insect biotypes resistant to Apparent Antassassin Insecticide and other Group 3A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect oppulation if Apparent Antassassin Insecticide or other Group 3A insecticide are used repeatedly. There was not effectiveness of Apparent Antassassin Insecticide or effectiveness of Apparent Antassassin Insecticide or esistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, AIRR Apparent Py Ltid accepts in liability for any losses that may result from the failure of Apparent Antassassin Insecticide to control section of the provided of the section of the provided of t Some naturally occurring insect biotypes resistant resistant insects. Apparent Antassassin Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, AIRR Apparent Ptv Ltd representative or local agricultural department agronomist. PROTECTION OF WILDLIFE, FISH,

## CRUSTACEANS AND ENVIRONMENT

Dangerous to fish and aquatic organisms. DO NOT contaminate dams, rivers, streams waterways or drains with product or the used container

#### PROTECTION OF PETS AND LIVESTOCK

Before treatment, remove animals and pets from the areas to be treated. Cover or remove any open food and water containers. Keep away from domestic pets, especially dogs. If dogs eat the granules, contact your in between payers, cracks, and crevices and dispersed into soil and/or lawn. Cover or remove fishponds. aquariums etc before application, DO NOT apply Apparent Antassassin Insecticide within 2 metres of a

This product is not classified as hazardous according to the Global Harmonization System (GHS) classification.

APVMA Approval No: 82922/130535 Batch No: Date of Manufacture: fishpond or water containing fish or aquatic species. **STORAGE AND DISPOSAL** 

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight.

Spillage - Dispose of waste as indicated below or according to the Australian Standard AS 2507 - Storage and Handling of Pesticides.

DO NOT allow spilled product to enter sewers, drains, creeks or any other waterways.

state or territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

# Wash hands after use.

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. SAFETY DATA SHEET (SDS)

Additional information is listed in the Safety Data Sheet which can be obtained from the supplier.
CONDITIONS OF SALE

The use of Apparent Antassassin Insecticide being beyond the control of the manufacturer no warranty expressed or implied is given by AIRR Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by veterinary surgeon immediately. Use with caution efficiency for any purpose for which it is used by around pets - ensure product is well dispersed i.e. swept the buyer, whether in accordance with the directions or not and AIRR Apparent Ptv Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.



READ SAFETY DIRECTIONS BEFORE OPENING OR USING



### AIRR APPARENT Pty Ltd

#### DIRECTIONS FOR USE RESTRAINTS:

DO NOT apply inside buildings.

SITUATIONS	vithin 2 m of water (p PEST	RATE (kg/ha)	RATE kg/100 m <sup>2</sup>	CRITICAL COMMENTS
Turf (For example: Lawns, commercial turf farms, parks, recreational areas, golf courses, bowling greens, sports fields)	Lawns Armyworm (Spodoptera maurita), Sod Webworm (Herpetogramma licarsisalis)	60	0.6	Broadcast Apparent Antassassin Insecticide with suitable application equipment to ensure uniform coverage over the treatment area. To ensure optimum control, irrigate the treated area with up to 4 mm of water immediately after application. Inspect treated areas for continuing activity. Re-apply as required.
	Argentine Stem Weevil adults ( <i>Listronotus</i> bonariensis)	60 - 120	0.6 – 1.2	
	African Black Beetle (Heteronychus arator)	120 - 180	1.2 – 1.8	
	Billbug adults (Spenophorus brunnipenis)	60 - 120	0.6 – 1.2	
	Black Ants, Coastal Brown Arits, Funnel Ants, Meat Ants, Sugar Ants, Stinging Ants only	60 - 220	0.6 – 2.2	Broadcast Apparent Antassassin Insecticide with suitable application equipment to ensure uniform coverage over the treatment area. Apply granules to areas where ants are active. Where possible apply granules directly to the nest. Use the low rate for maintenance treatments or to control light infestations. Use the high rate for heavy infestations. Use the high rate for heavy infestations and for maximum residual control. The elimination of funnel ants from a particular site will generally require more than one application. Initial applications should be broadcast over affected areas. As the number of active colonies is reduced, application should shift to targeting active mounds.
External surrounds of building and structures: Gardens, Lawns, Ant nests & trails	Black Ants, Coastal Brown Ants, Funnel Ants, Meat Ants, Sugar Ants, Stinging Ants only	60 - 220	0.6 – 2.2	Apply granules to areas where ants are active. Where possible apply granules directly to the nest. Use the low rate for maintenance treatments or to control light infestations. Use the high rate for heavy infestations and for maximum residual control. Perimeter Application. Where perimeter pest control is being implemented apply Apparent Antassassin insecticide in a two or three metre band, around and adjacent to the building or structure.  Pad Mounted Electrical Sites: To reduce or prevent ant mound building activity on electrical pads, uniformly distribute 60 to 220 grams of Apparent Antassassin insecticide per 10 m² of mounting pad. Where existing transformers are completely or partially filled with soil excavated by resident ants, remove any excess fill and level the remaining soll prior to apolication.